

You've identified a patient in your practice who may have uncontrolled gout

PRINT OUT THIS PACKET AND FILL OUT THE FORMS TO GET YOUR PATIENT REFERRED TO A GOUT SPECIALIST



CHECKLIST

- Inform the patient they are being referred to a gout specialist
- Give your patient the printed Patient Guide
- Fill out the Referral Support Form, and ask your patient to sign it so they can get connected to a Gout Nurse Advocate
- Submit the completed Referral Support Form by fax or email
- Follow your office's standard referral process to refer the patient to a gout specialist
- Send any patient history and lab work to the uncontrolled gout specialist's office, if applicable

If you have any questions, call your Amgen representative.

You've been referred to a gout specialist

WHAT'S NEXT?

Your doctor believes you may have uncontrolled gout. A gout specialist can help.



Connect with a Gout Nurse Advocate (GNA)* to prepare for your upcoming visit. Your GNA will:

- Provide information about uncontrolled gout
- Explain how KRYSTEXXA may be able to help

Simply complete the Referral Support Form or call 1-833-469-4688 to speak with a GNA



KRYSTEXXA is the first and only FDA-approved treatment for uncontrolled gout

- Reduces nonresolving tophi (or gouty lumps), high uric acid levels, and gout flares over time
- Works differently than oral medication
- Starts working within 24 hours of the first treatment



Follow up with your gout specialist directly if they haven't contacted you within 1 to 2 weeks

Dr: _____ Phone Number: _____

*GNAs do not provide individual patient care or medical advice.



Take action today by preparing for your specialist visit

Point your smartphone camera at this QR code or visit [PrepareForYourVisit.com](https://www.prepareforyourvisit.com) to personalize your doctor discussion guide. The more you share with them, the better they can help.

SELECT IMPORTANT SAFETY INFORMATION

What is the most important information I should know about KRYSTEXXA?

Serious allergic reactions may happen in some patients who receive KRYSTEXXA. These allergic reactions can be life-threatening and usually happen within 2 hours of the infusion.

KRYSTEXXA should be given to you by a doctor or nurse in a healthcare setting where serious allergic reactions can be treated. Your doctor or nurse should watch you for any signs of a serious allergic reaction during and after your treatment with KRYSTEXXA.

Please see additional Important Safety Information on back and Medication Guide at [KRYSTEXXA.com](https://www.krystexxa.com).

KRYSTEXXA
pegloticase

USE

KRYSTEXXA® (pegloticase) is a prescription medicine used in adults to help reduce the signs and symptoms of gout that are not controlled by other treatments.

KRYSTEXXA is not for use in people with too much uric acid in their bodies who do not have symptoms (asymptomatic hyperuricemia).

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about KRYSTEXXA?

Serious allergic reactions may happen in some patients who receive KRYSTEXXA. These allergic reactions can be life-threatening and usually happen within 2 hours of the infusion.

KRYSTEXXA should be given to you by a doctor or nurse in a healthcare setting where serious allergic reactions can be treated. Your doctor or nurse should watch you for any signs of a serious allergic reaction during and after your treatment with KRYSTEXXA.

Tell your doctor or nurse right away if you have any of these symptoms during or after your treatment with KRYSTEXXA:

- wheezing, shortness of breath, cough, chest tightness, chest pain, or trouble breathing
- dizziness, fainting, fast or weak heartbeat or feeling nervous
- reddening of the face, itching, hives, or feeling warm
- swelling of the throat or tongue, throat tightness, hoarse voice, or trouble swallowing

Who should not receive KRYSTEXXA?

Do not receive KRYSTEXXA if you:

- have a rare blood problem called glucose-6-phosphate dehydrogenase (G6PD) deficiency or favism. Your doctor may test you for G6PD before you start KRYSTEXXA.
- have had a serious allergic reaction to KRYSTEXXA or any of its ingredients. See the end of the Medication Guide for a complete list of ingredients in KRYSTEXXA.

What should I tell my doctor before receiving treatment with KRYSTEXXA?

Before you receive KRYSTEXXA, tell your doctor about all of your medical conditions, including if you:

- ever had any heart problems or high blood pressure.
- are pregnant or plan to become pregnant. It is not known if KRYSTEXXA will harm your unborn baby. Talk to your doctor if you are pregnant or plan to become pregnant.
- are breastfeeding or plan to breastfeed. It is not known if KRYSTEXXA passes into your breast milk. You and your doctor should decide if you will receive KRYSTEXXA or breastfeed.

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. **Do not** take any other uric acid lowering drug, such as allopurinol, febuxostat (Uloric), or probenecid, while receiving KRYSTEXXA.

KRYSTEXXA is recommended to be given with another prescription medicine called methotrexate. KRYSTEXXA may also be used alone. You and your doctor will decide the treatment that is right for you.

Prior to your treatment with KRYSTEXXA, your doctor may give you medicine to help reduce your risk of getting gout flares or an allergic reaction. Take these medicines as directed by your doctor or nurse. Your doctor will also test your uric acid levels prior to each treatment to monitor your response to KRYSTEXXA.

Your gout flares may increase in the first 3 months when you start receiving KRYSTEXXA. It's important to understand that this is happening because KRYSTEXXA is breaking down uric acid in your body. **Do not** stop receiving KRYSTEXXA even if you have a flare, as the amount of flares will decrease after 3 months of treatment. Your doctor may give you other medicines to help reduce your gout flares for the first few months after starting KRYSTEXXA.

What are the possible side effects of KRYSTEXXA?

In KRYSTEXXA clinical trials:

The most common side effects of KRYSTEXXA when given together with methotrexate were gout flares, joint pain, coronavirus disease 2019 (COVID-19), nausea, and fatigue.

The most common side effects of KRYSTEXXA were gout flares, allergic reactions (including infusion reactions). See "What is the most important information I should know about KRYSTEXXA?", nausea, bruising, sore throat, constipation, chest pain, coronavirus disease 2019 (COVID-19), and vomiting.

This is not a complete list of all possible side effects. Call your doctor for medical advice about side effects.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/safety/medwatch or call 1-800-FDA-1088.

For additional Important Safety Information, please see the Medication Guide at KRYSTEXXA.com and discuss with your doctor.